



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION OF PROBLEM 226.

BY PROF. W. W. JOHNSON, ST. JOHN'S COLLEGE, ANNAPOLIS, MD.

PUTTING $\frac{2k\pi}{n} = \theta$, $x = \cos \theta$ in the first of the required equations, and $x = \sin \theta$ in the second. In either case, $\sin n\theta = 0$, and $\cos n\theta = 1$; hence if we can express either of these quantities in a finite number of terms involving powers of $\cos \theta$ we shall have the first of the required equations, and if we express either of them in terms of $\sin \theta$, we have the second equation. Now $\sin n\theta$ cannot be expressed in a finite number of terms involving $\cos \theta$; but for $\cos n\theta$ we have (See Chauvenet's Trig., eq. 460, page 138),

$$\begin{aligned} \cos n\theta = \cos \frac{n\pi}{2} & \left[1 - \frac{n^2}{2} \cos^2 \theta + \frac{n^2(n^2-2^2)}{1.2.3.4} \cos^4 \theta + \dots \right] \\ & + \sin \frac{n\pi}{2} \left[n \cos \theta - \frac{n(n^2-1^2)}{1.2.3} \cos^3 \theta + \frac{n(n^2-1^2)(n^2-3^2)}{5!} \cos^5 \theta \dots \right]. \end{aligned}$$

Substituting $\cos \theta = x$ we have, when n is an odd integer,

$$\begin{aligned} 0 = -1 + (-1)^{\frac{n-1}{2}} & \left[nx - \frac{n(n^2-1^2)}{1.2.3} x^3 + \frac{n(n^2-1^2)(n^2-3^2)}{5!} x^5 - \dots \right. \\ & \left. + (-1)^{\frac{n-1}{2}} \frac{n(n^2-1^2) \dots [n^2-(n-2)^2]}{n!} x^n \right], \quad (1) \end{aligned}$$

but when n is even we have

$$\begin{aligned} 0 = -1 + (-1)^{\frac{n}{2}} & \left[1 - \frac{n^2}{2} x^2 + \frac{n^2(n^2-2^2)}{4!} x^4 - \dots \right. \\ & \left. + (-1)^{\frac{n}{2}} \frac{n^2(n^2-2^2) \dots [n^2-(n-2)^2]}{n!} x^n \right]. \quad (2) \end{aligned}$$

Equations (1) and (2) are therefore the equations whose roots are

$$x = \cos \frac{2k\pi}{n}.$$

For the second equation $x = \sin \theta$, and we are required to express either $\cos n\theta$ or $\sin n\theta$ in terms of $\sin \theta$. We have

$$\begin{aligned} \cos n\theta &= 1 - \frac{n^2}{2} \sin^2 \theta + \frac{n^2(n^2-2^2)}{1.2.3.4} \sin^4 \theta - \frac{n^2(n^2-2^2)(n^2-4^2)}{6!} \sin^6 \theta + \dots \\ \sin n\theta &= n \sin \theta - \frac{n(n^2-1^2)}{1.2.3} \sin^3 \theta + \frac{n(n^2-1^2)(n^2-3^2)}{5!} \sin^5 \theta - \dots \end{aligned}$$

When n is odd, the expression for $\sin n\theta$ terminates, and we have by substitution

$$0 = x - \frac{n^2-1^2}{1.2.3} x^3 + \frac{(n^2-1^2)(n^2-3^2)}{5!} x^5 - \dots \\ + (-1)^{\frac{n-1}{2}} \frac{(n^2-1^2)(n^2-3^2) \dots [n^2-(n-2)^2]}{n!} x^n, \quad (3)$$

but when n is even the expression for $\cos \theta$ terminates, and we have, on substitution,

$$0 = x^2 - \frac{n^2-2^2}{3.4} x^4 + \frac{(n^2-2^2)(n^2-4^2)}{3.4.5.6} x^6 - \dots \\ - (-1)^{\frac{n}{2}} \frac{(n^2-2^2) \dots [n^2-(n-2)^2]}{3.4.5 \dots n} x^n. \quad (4)$$

Equations (3) and (4) are therefore those whose roots are

$$x = \sin \frac{2k\pi}{n}.$$

The several roots are in each case found by giving to k the values $0, 1, 2, \dots, n-1$. Putting $k=0$ we see that $x=1$ is a root of equations (1) and (2), whence we have the numerical series, when n is odd,

$$(-1)^{\frac{n-1}{2}} = n - \frac{n(n^2-1^2)}{1.2.3} + \frac{n(n^2-1^2)(n^2-3^2)}{5!} - \dots$$

and when n is even,

$$(-1)^{\frac{n}{2}} = 1 - \frac{n^2}{2} + \frac{n^2(n^2-2^2)}{4!} - \frac{n^2(n^2-2^2)(n^2-4^2)}{6!} + \dots$$

Since $\cos \frac{2k\pi}{n} = \cos \frac{2(n-k)\pi}{n}$ the roots (exclusive of unity) of eq. (1) occur in equal pairs, and the equation must be of the form

$$(x-1)[F_1(x)]^2 = 0,$$

and for a like reason equation (2) must have the form

$$(x^2-1)[F_2(x)]^2 = 0.$$

The equations $F_1(x) = 0$ and $F_2(x) = 0$ are therefore those whose roots are the *different* values of x , exclusive of 1 and -1 , which are represented by $\cos(2k\pi \div n)$.

I proceed to find the form of the functions $F_1(x)$ and $F_2(x)$. Observe in the first place that we may, in any one of the equations, change n to $-n$, in each of the coefficients; and also that, since in (1) and (2) the degree of each term in n is the same as its degree in x , the same will be true in the quotients when we divide these equations by $-1+x$ or $-1+x^2$, and also in the square roots of the results, that is in the functions F_1 and F_2 .

Assuming n to have the form $4n'+1$ in eq. (1) we may give to n the successive values $1, -3, 5, -7, 9$ etc., and by changing the sign of n we shall

have the form which the function takes when n is of the form $4n' + 3$, which form may be denoted by F_3 , the admissible values of n being $-1, 3, -5, 7$ etc. It is obvious that the first term of $F_1(x)$ is 1; and, since the expression must reduce to its first term when $n = 1$, every term after the first must contain the factor $n-1$. In like manner all terms after the second must contain the factor $n + 3$, since when $n = -3$ all terms after the second must vanish; and so on. Thus we may assume

$$F_1(x) = 1 + A(n-1)x + B(n-1)(n+3)x^2 + \dots$$

in which A, B , etc., are numerical factors, since the degree of each term in n is to be the same as its degree in x . Then the expression $F_3(x)$, found by changing n to $-n$, is

$$F_3(x) = 1 - A(n+1)x + B(n+1)(n-3)x^2 - \dots$$

In either case the last term contains $x^{\frac{n-1}{2}}$, and $\frac{n-1}{2}$ factors which are successive multiples of 4, the least being 4; hence, denoting the numerical factor by L , the value of this term is

$$L \cdot 4^{\frac{n-1}{2}} \left(\frac{n-1}{2}\right)! x^{\frac{n-1}{2}}.$$

Now this term is the square root of the last term of $[F_1(x)]^2$ which is the result of dividing the second member of (1) by $-1+x$. The latter term may be written in the form

$$\frac{(2n-2)(2n-4)\dots(n+1)(n-1)\dots 2}{(n-1)!} x^{n-1},$$

$$= 2^{n-1} x^{n-1},$$

the square root of which is $\pm 2^{\frac{n-1}{2}} x^{\frac{n-1}{2}}$. Theorefore

$$L \cdot 4^{\frac{n-1}{2}} \left(\frac{n-1}{2}\right)! x^{\frac{n-1}{2}} = \pm 2^{\frac{n-1}{2}} x^{\frac{n-1}{2}},$$

whence

$$L = \pm \frac{1}{2^{\frac{n-1}{2}} \left(\frac{n-1}{2}\right)!},$$

which gives the value of each coefficient, since any term may be made the last term in one or other of the expressions $F_1(x)$ and $F_3(x)$. To determine the signs of the terms, observe that the coefficient of the term preceding the last in $[F_1(x)]^2$ is the same as that of the last term, and is therefore positive; hence the last two terms of $F_1(x)$ or $F_3(x)$ have the same sign. Then since $F_1(x)$ contains an odd number of terms A and B must have the same sign, and so must C and D, E and F , etc., and since $F_3(x)$ contains an even number of terms, 1 and $-A, B$ and $-C$, etc., must have the same

sign. Thus we have when $n = 4n' + 1$

$$F_1(x) = 1 - \frac{n-1}{2}x - \frac{(n-1)(n+3)}{2^2 \cdot 2!}x^2 + \frac{(n-1)(n+3)(n-5)}{2^3 \cdot 3!}x^3 + \dots$$

and when $n = 4n' + 3$

$$F_3(x) = 1 + \frac{n+1}{2}x - \frac{(n+1)(n-3)}{2^2 \cdot 2!}x^2 - \frac{(n+1)(n-3)(n+5)}{2^3 \cdot 3!}x^3 + \dots$$

Eq. (2) takes different forms when n is of the form $4n' + 2$, and when it is of the form $4n'$. In the first case, all the pairs of equal roots occur in couples having numerically equal positive and negative values, so that the equation may be written in the form $(x^2 - 1)[F_2(x^2)]^2 = 0$; but in the second case there is a single pair of zero roots, so that the form of the equation is $x^2(x^2 - 1)[F_4(x^2)]^2 = 0$.

When $n = 4n' + 2$, eq. (2) becomes

$$0 = -1 + \frac{n^2}{2 \cdot 2}x^2 - \frac{n^2(n^2 - 2^2)}{2 \cdot 4!}x^4 + \dots + \frac{n^2(n^2 - 2^2) \dots [n^2 - (n-2)^2]}{2 \cdot n!}x^n$$

in which n admits of the values $\pm 2, \pm 6, \pm 10$, etc. The first term of $F_2(x^2)$ is unity; and, since the expression must reduce to its first term when $n = \pm 2$, all the terms beyond the first must contain $n^2 - 2^2$ as a factor. In like manner, all the terms beyond the second must contain $n^2 - 6^2$, etc. Thus we may assume

$$F_2(x^2) = 1 + A(n^2 - 2^2)x^2 + B(n^2 - 2^2)(n^2 - 6^2)x^4 + \dots + L(n^2 - 2^2) \dots \times [n^2 - (n-4)^2]x^{\frac{n-2}{2}},$$

in which A, B , etc., have numerical values. The last term may be written in the form

$$L \cdot (2n-4)(2n-8) \dots (n+2)(n-2) \dots 4x^{\frac{n-2}{2}} = L \cdot 4^{\frac{n-2}{2}} \left(\frac{n-2}{2}\right)! x^{\frac{n-2}{2}}.$$

But the last term of the quotient, when the above form of eq. (2) is divided by $-1 + x^2$, is

$$\frac{(2n-2)(2n-4) \dots (n+2)n(n-2) \dots 2}{2 \cdot (n-1)!} x^{n-2} = 2^{n-2} x^{n-2},$$

the square root of which is $\pm 2^{\frac{n-2}{2}} x^{\frac{n-2}{2}}$. Hence

$$L \cdot 4^{\frac{n-2}{2}} \left(\frac{n-2}{2}\right)! x^{\frac{n-2}{2}} = \pm 2^{\frac{n-2}{2}} x^{\frac{n-2}{2}},$$

and

$$L = \pm \frac{1}{2^{\frac{n-2}{2}} \left(\frac{n-2}{2}\right)!},$$

which gives the form of the numerical factors. It is easily shown that the last term but one of $F_2(x^2)$ is negative, hence the last two terms of $F_2(x^2)$ are always of opposite signs, and we have

$$F_2(x^2) = 1 - \frac{n^2-2^2}{2^2 \cdot 2!} x^2 + \frac{(n^2-2^2)(n^2-6^2)}{2^4 \cdot 4!} x^4 - \dots$$

When n is of the form $4n'$, eq. (2) becomes

$$0 = -x^2 + \frac{n^2-2^2}{3 \cdot 4} x^4 - \frac{(n^2-2^2)(n^2-4^2)}{3 \cdot 4 \cdot 5 \cdot 6} x^6 + \dots + \frac{2(n^2-2^2) \dots [n^2-(n-2)^2]}{n!} x^n,$$

in which n admits of the values $\pm 4, \pm 8$, etc., the form of the equation being $x^2(x^2-1)[F_4(x^2)]^2 = 0$, so that the degree of each term of $F_4(x^2)$ in n will still be the same as the degree in x . The first term of $F_4(x^2)$ is unity, and it may be assumed in the form

$$F_4(x^2) = 1 + A(n^2-4^2)x^2 + B(n^2-4^2)(n^2-8^2)x^4 + \dots + L(n^2-4^2) \dots \times [n^2-(n-4)^2] x^{\frac{n-4}{2}},$$

A, B , etc., having numerical values. In this case, the last term is

$$L \cdot (2n-4)(2n-8) \dots (n+4)(n-4) \dots 4 \cdot x^{\frac{n-4}{2}} = L \frac{4^{\frac{n-2}{2}} \left(\frac{n-2}{2}\right)!}{n} x^{\frac{n-4}{2}}.$$

But the last term of $[F_4(x)]^2$ may be written

$$\frac{2(2n-2)(2n-4) \dots (n+2)n(n-2) \dots 2}{n \cdot n!} x^{n-4} \\ = \frac{2^n(n-1)!}{n^2(n-1)!} x^{n-4} = \frac{2^n}{n^2} x^{n-4},$$

the square root of which is $\pm \frac{2^{\frac{1}{2}n} x^{\frac{n-4}{2}}}{n}$. Hence

$$L \frac{4^{\frac{n-2}{2}} \left(\frac{n-2}{2}\right)!}{n} x^{\frac{n-4}{2}} = \pm \frac{2^{\frac{1}{2}n} x^{\frac{n-4}{2}}}{n}, \text{ and } L = \pm \frac{1}{2^{\frac{n-4}{2}} \left(\frac{n-2}{2}\right)!}.$$

The terms are of alternate signs, for the reason already given in the case of F_2 , hence we have

$$F_4(x^2) = 1 - \frac{n^2-4^2}{2^2 \cdot 3!} x^2 + \frac{(n^2-4^2)(n^2-8^2)}{2^4 \cdot 5!} x^4 - \dots$$

Of equation (3) we can only infer, what is obvious on inspection, that it takes the form $xF(x^2) = 0$; but eq. (4) takes the form $x^2[F(x^2)]^2 = 0$ when n has the form $4n'+2$, and the form $x^2(x^2-1)[F(x^2)]^2 = 0$, when n has the form $4n'$. In the latter case, the equation becomes identical with eq. (2) in the like case, and in the former case, we find by the method pursued above

$$F(x^2) = 1 - \frac{n^2-2^2}{2^2 \cdot 3!} x^2 + \frac{(n^2-2^2)(n^2-6^2)}{2^4 \cdot 5!} x^4 - \dots$$